

Anti-Human 4-1BB Ligand Antibody
Catalog # ABG10001**Specification**

Anti-Human 4-1BB Ligand Antibody - Product Information

Application	WB, IHC, E
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

Anti-Human 4-1BB Ligand Antibody - Additional Information**Preparation**

Produced from sera of rabbits pre-immunized with highly pure recombinant Human 4-1BBL. Anti-Human 4-1BBL specific antibody was purified by affinity chromatography employing immobilized Human 4-1BBL matrix.

WesternBlot

To detect h4-1BBL by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. When used in conjunction with compatible secondary reagents, the detection limit for recombinant h4-1BBL is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

Sandwich

To detect h4-1BBL by sandwich ELISA (using 100µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems's Biotinylated Anti-Human 4-1BBL (60-068BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant h4-1BBL.

Immunohistochemistry

This antibody stained formalin-fixed, paraffin-embedded sections of human colon/rectum adenocarcinoma. The recommended concentration is 0.25 mg/ml with an overnight incubation at 4°C. An HRP-labeled polymer detection system was used with a DAB chromogen. Proteinase K antigen retrieval is recommended. Optimal concentrations and conditions may vary.

Formulation

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Storage

-20°C

Precautions

Anti-Human 4-1BB Ligand Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-Human 4-1BB Ligand Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Human 4-1BB Ligand Antibody - Images