

TRIM11 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP13284B**Specification**

TRIM11 Antibody (C-term) - Product Information

| | |
|-------------------|---|
| Application | WB, IHC-P,E |
| Primary Accession | Q96F44 |
| Other Accession | B1H278 , A0JN74 , NP_660215.1 |
| Reactivity | Human |
| Predicted | Bovine, Rat |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | Rabbit IgG |
| Calculated MW | 52774 |
| Antigen Region | 300-329 |

TRIM11 Antibody (C-term) - Additional Information**Gene ID** 81559**Other Names**

E3 ubiquitin-protein ligase TRIM11, 632-, Protein BIA1, RING finger protein 92, Tripartite motif-containing protein 11, TRIM11, RNF92

Target/Specificity

This TRIM11 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 300-329 amino acids from the C-terminal region of human TRIM11.

Dilution

WB~~1:1000
IHC-P~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

TRIM11 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

TRIM11 Antibody (C-term) - Protein Information**Name** TRIM11 {ECO:0000303|PubMed:16904669, ECO:0000312|HGNC:HGNC:16281}

Function E3 ubiquitin-protein ligase that promotes the degradation of insoluble ubiquitinated proteins, including insoluble PAX6, poly-Gln repeat expanded HTT and poly-Ala repeat expanded ARX (By similarity). Mediates PAX6 ubiquitination leading to proteasomal degradation, thereby modulating cortical neurogenesis (By similarity). May also inhibit PAX6 transcriptional activity, possibly in part by preventing the binding of PAX6 to its consensus sequences (By similarity). May contribute to the regulation of the intracellular level of HN (humanin) or HN-containing proteins through the proteasomal degradation pathway (By similarity). Mediates MED15 ubiquitination leading to proteasomal degradation (PubMed:[16904669](#)). May contribute to the innate restriction of retroviruses (PubMed:[18248090](#)). Upon overexpression, reduces HIV-1 and murine leukemia virus infectivity, by suppressing viral gene expression (PubMed:[18248090](#)). Antiviral activity depends on a functional E3 ubiquitin-protein ligase domain (PubMed:[18248090](#)). May regulate TRIM5 turnover via the proteasome pathway, thus counteracting the TRIM5-mediated cross-species restriction of retroviral infection at early stages of the retroviral life cycle (PubMed:[18248090](#)). Acts as an inhibitor of the AIM2 inflammasome by promoting autophagy-dependent degradation of AIM2 (PubMed:[27498865](#)). Mechanistically, undergoes autoubiquitination upon DNA stimulation, promoting interaction with AIM2 and SQSTM1/p62, leading to AIM2 recruitment to autophagosomes (PubMed:[27498865](#)).

Cellular Location

Cytoplasm. Nucleus


Tissue Location

Ubiquitous..

TRIM11 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TRIM11 Antibody (C-term) - Images

TRIM11 Antibody (C-term) (Cat. #AP13284b) western blot analysis in HepG2 cell line lysates

(35ug/lane). This demonstrates the TRIM11 antibody detected the TRIM11 protein (arrow).



TRIM11 Antibody (C-term) (Cat. #AP13284b) immunohistochemistry analysis in formalin fixed and paraffin embedded human heart tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of TRIM11 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

TRIM11 Antibody (C-term) - Background

The protein encoded by this gene is a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. This protein localizes to the nucleus and the cytoplasm. Its function has not been identified. [provided by RefSeq].

TRIM11 Antibody (C-term) - References

Aurino, S., et al. Acta Myol 27, 90-97 (2008) :
Hong, S.J., et al. Biochem. Biophys. Res. Commun. 368(3):650-655(2008)
Uchil, P.D., et al. PLoS Pathog. 4 (2), E16 (2008) :
Ishikawa, H., et al. FEBS Lett. 580(20):4784-4792(2006)
Cooper, S.T., et al. BMC Genet. 6, 43 (2005) :

TRIM11 Antibody (C-term) - Citations

- [Cutaneous Melanocytoma With CRTC1-TRIM11 Fusion: Report of 5 Cases Resembling Clear Cell Sarcoma.](#)