

## SMO Antibody (Center)

Peptide Affinity Purified Rabbit Polyclonal Antibody (Pab)  
Catalog # AP16325c

### Specification

#### SMO Antibody (Center) - Product Information

Application	WB,E
Primary Accession	<a href="#">O99835</a>
Other Accession	<a href="#">NP_005622.1</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	86397
Antigen Region	539-567

#### SMO Antibody (Center) - Additional Information

Gene ID 6608

#### Other Names

Smoothened homolog, SMO, Protein Gx, SMO, SMOH

#### Target/Specificity

This SMO antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 539-567 amino acids from the Central region of human SMO.

#### Dilution

WB~~1:1000

#### Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.


#### Precautions

SMO Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

#### SMO Antibody (Center) - Protein Information

Name SMO

Synonyms SMOH



#### SMO Antibody (Center) - Background

The protein encoded by this gene is a G protein-coupled receptor that interacts with the patched protein, a receptor for hedgehog proteins. The encoded protein transduces signals to other proteins after activation by a hedgehog protein/patched protein complex.

#### SMO Antibody (Center) - References

- Zhang, L., et al. Oral Dis 16(8):818-822(2010)
- Desch, P., et al. Oncogene 29(35):4885-4895(2010)
- Walter, K., et al. Clin. Cancer Res. 16(6):1781-1789(2010)
- Hirotsu, M., et al. Mol. Cancer 9, 5 (2010) :
- Rittie, L., et al. Aging Cell 8(6):738-751(2009)

**Function**

G protein-coupled receptor that probably associates with the patched protein (PTCH) to transduce the hedgehog's proteins signal. Binding of sonic hedgehog (SHH) to its receptor patched is thought to prevent normal inhibition by patched of smoothed (SMO). Required for the accumulation of KIF7, GLI2 and GLI3 in the cilia (PubMed:<a href="http://www.uniprot.org/citations/19592253" target="\_blank">19592253</a>). Interacts with DLG5 at the ciliary base to induce the accumulation of KIF7 and GLI2 at the ciliary tip for GLI2 activation (By similarity).

**Cellular Location**

Membrane; Multi-pass membrane protein.  
Cell projection, cilium

**SMO Antibody (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**SMO Antibody (Center) - Citations**

- [The sonic hedgehog pathway mediates Tongxinluo capsule-induced protection against blood-brain barrier disruption after ischemic stroke in mice.](#)
- [Dehydroeffusol inhibits viability and epithelial-mesenchymal transition through the Hedgehog and Akt/mTOR signaling pathways in neuroblastoma cells.](#)
- [Salinomycin exerts anticancer effects on human breast carcinoma MCF-7 cancer stem cells via modulation of Hedgehog signaling.](#)