

**PACSIN2 Antibody (C-term)**  
Purified Rabbit Polyclonal Antibody (Pab)  
Catalog # AP8088b

**Specification**

**PACSIN2 Antibody (C-term) - Product Information**

Application	IHC-P, WB,E
Primary Accession	<a href="#">Q9UNF0</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Antigen Region	342-371

**PACSIN2 Antibody (C-term) - Additional Information**

Gene ID 11252

**Other Names**

Protein kinase C and casein kinase substrate in neurons protein 2, Syndapin-2, Syndapin-II, PACSIN2

**Target/Specificity**

This PACSIN2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 342-371 amino acids from the C-terminal region of human PACSIN2.

**Dilution**

IHC-P ~ 1:50~100  
WB ~ 1:1000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**


PACSIN2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**PACSIN2 Antibody (C-term) - Protein Information**

Name PACSIN2



Immunohistochemical analysis of paraffin-embedded H. stomach section using PACSIN2 Antibody (C-term) (Cat#AP8088b). AP8088b was diluted at 1:25 dilution. A peroxidase-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody, followed by DAB staining.



PACSIN2 Antibody (S357) (Cat.# AP8088b) western blot analysis in Daudi, HL-60 cell line and mouse brain, rat heart lysates (35ug/lane). This demonstrates the PACSIN2 antibody detected the PACSIN2 protein (arrow).

## Function


Lipid-binding protein that is able to promote the tubulation of the phosphatidic acid-containing membranes it preferentially binds. Plays a role in intracellular vesicle-mediated transport. Involved in the endocytosis of cell-surface receptors like the EGF receptor, contributing to its internalization in the absence of EGF stimulus. May also play a role in the formation of caveolae at the cell membrane. Recruits DNMT2 to caveolae, and thereby plays a role in caveola-mediated endocytosis.

## Cellular Location

Cytoplasm. Cytoplasm, cytoskeleton. Cytoplasmic vesicle membrane; Peripheral membrane protein; Cytoplasmic side. Early endosome Recycling endosome membrane. Cell projection, ruffle membrane; Peripheral membrane protein; Cytoplasmic side. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection. Membrane, caveola. Note=Detected at the neck of flask-shaped caveolae. Localization to tubular recycling endosomes probably requires interaction with MICALL1 and EHD1

## Tissue Location

Widely expressed.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

## PACSIN2 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## PACSIN2 Antibody (C-term) - Citations

- [A disease-associated frameshift mutation in caveolin-1 disrupts caveolae formation and function through introduction of a de novo ER retention signal.](#)
- [Clostridium difficile Toxin A Undergoes Clathrin-Independent, PACSIN2-Dependent Endocytosis.](#)
- [Characterization of a caveolin-1 mutation associated with both PAH and congenital generalized lipodystrophy.](#)
- [EHD3 is Required for Tubular Recycling Endosome Stabilization and an Asparagine-Glutamic Acid Residue Pair within its EH Domain Dictates its Selective Binding to NPF Peptides.](#)
- [Endocytic recycling protein EHD1 regulates primary cilia morphogenesis and SHH signaling during neural tube development.](#)
- [Differential roles of C-terminal Eps15 homology domain proteins as vesiculators and tubulators of recycling endosomes.](#)
- [Cooperation of MICAL-L1, syndapin2, and phosphatidic acid in tubular recycling endosome biogenesis.](#)
- [The F-BAR protein PACSIN2 regulates epidermal growth factor receptor internalization.](#)

## PACSIN2 Antibody (C-term) - Background

PACSIN may play a role in vesicle formation and transport. This protein homo- and hetero-aggregates with other PACSINs. It also binds dynamin 1, synaptojanin, synapsin 1 and the neural Wiskott-Aldrich syndrome protein (N-WASP). The protein exhibits a vesicle-like cytoplasmic distribution and is ubiquitously expressed. PACSIN is phosphorylated by casein kinase 2 (CK2) and protein kinase C (PKC). The protein contains 1 FCH domain and 1 SH3 domain.

## PACSIN2 Antibody (C-term) - References

Strausberg, R.L., et al., Proc. Natl. Acad. Sci. U.S.A. 99(26):16899-16903 (2002). Wiemann, S., et al., Genome Res. 11(3):422-435 (2001). Ritter, B., et al., FEBS Lett. 454(3):356-362 (1999). Dunham, I., et al., Nature 402(6761):489-495 (1999).

- [Pacsin 2 is recruited to caveolae and functions in caveolar biogenesis.](#)
- [The F-BAR domain protein PACSIN2 associates with Rac1 and regulates cell spreading and migration.](#)