

Anti-Bovine IgG F(c) Secondary Antibody
Rabbit Polyclonal, Unconjugated
Catalog # ASR1751**Specification****Anti-Bovine IgG F(c) Secondary Antibody - Product Information**

Description	Anti-BOVINE IgG F(c) (RABBIT) Antibody
Host	Rabbit
Conjugate	Unconjugated
Target Species	Bovine
Reactivity	Bovine
Clonality	Polyclonal
Application	,1,10,15,
Application Note	ELISA 1:20,000-1:100,000;Western Blot 1:2,000-1:10,000;Immunochemistry 1:1,000-1:5,000
Physical State	Liquid (sterile filtered)
Host Isotype	IgG
Target Isotype	IgG F(c)
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Anti-Bovine IgG F(c) was produced by repeated immunization with bovine IgG F(c) fragment in rabbit.
Stabilizer	None
Preservative	0.01% (w/v) Sodium Azide

Anti-Bovine IgG F(c) Secondary Antibody - Additional Information**Shipping Condition**

Wet Ice

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Bovine IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Bovine IgG, Bovine IgG F(c) and Bovine Serum. No reaction was observed against Bovine IgG F(ab')₂.

Storage Condition

Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Bovine IgG F(c) Secondary Antibody - Protein Information

Anti-Bovine IgG F(c) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Bovine IgG F(c) Secondary Antibody - Images

Anti-Bovine IgG F(c) Secondary Antibody - Background

Anti-Bovine IgG F(c) antibody generated in rabbit detects specifically bovine IgG. This secondary antibody anti-Bovine is ideal for investigators who routinely perform titration assays, western-blot, immunoprecipitation and more generally immunoassays.