

SOX2 Antibody
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AM2048a

Specification

SOX2 Antibody - Product Information

Application	IF, WB, IHC-P, FC,E
Primary Accession	P48431
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1

SOX2 Antibody - Additional Information

Gene ID 6657

Other Names

Transcription factor SOX-2, SOX2

Target/Specificity

SOX2 recombinant protein is used to produce this monoclonal antibody.

Dilution

IF~~1:10~50
WB~~1:4000
IHC-P~~1:50~100
FC~~1:10~50

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions


SOX2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

SOX2 Antibody - Protein Information


Name SOX2

Function

Transcription factor that forms a trimeric complex with OCT4 on DNA and controls the expression of a number of genes involved in embryonic development such as YES1, FGF4, UTF1 and ZFP206 (By similarity). Binds to the



Fluorescent confocal image of SY5Y cells stained with SOX2 antibody. SY5Y cells were fixed with 4% PFA (20 min), permeabilized with Triton X-100 (0.2%, 30 min). Cells were then incubated with AM2048a SOX2 primary antibody (1:100, 2 h at room temperature). For secondary antibody, Alexa Fluor® 488 conjugated donkey anti-mouse antibody (green) was used (1:1000, 1h). Note the highly specific localization of the SOX2 mainly to the nucleus.



Fluorescent image of A549 cell stained with SOX2 Antibody(Cat#AM2048a/SG110310AA).A549 cells were fixed with 4% PFA (20 min),

proximal enhancer region of NANOG (By similarity). Critical for early embryogenesis and for embryonic stem cell pluripotency (PubMed: 18035408). Downstream SRRF target that mediates the promotion of neural stem cell self-renewal (By similarity). Keeps neural cells undifferentiated by counteracting the activity of proneural proteins and suppresses neuronal differentiation (By similarity). May function as a switch in neuronal development (By similarity).

Cellular Location


Nucleus {ECO:0000250|UniProtKB:P48432}.

SOX2 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


permeabilized with Triton X-100 (0.1%, 10 min), then incubated with SOX2 primary antibody (1:25, 1 h at 37°C. For secondary antibody, Alexa Fluor® 488 conjugated donkey anti-mouse antibody (green) was used (1:400, 50 min at 37°C. Cytoplasmic actin was counterstained with Alexa Fluor® 555 (red) conjugated Phalloidin (7 units/ml, 1 h at 37°C. SOX2 immunoreactivity is localized to Nucleus significantly.




Western blot analysis of SOX2 (arrow) using mouse monoclonal SOX2 antibody (Cat.#AM2048a). 293 cell lysates (2 µg/lane) either nontransfected (Lane 1) or transiently transfected with the SOX2 gene (Lane 2) (Origene Technologies)




Western blot analysis of SOX2 Antibody (Cat.#AM2048a) by SOX2 recombinant protein. SOX2 (arrow) was detected using the purified Mab.



Western blot analysis of lysate from NCCIT cell line, using SOX2 Antibody (Cat. #AM2048a). AM2048a was diluted at 1:1000. A goat anti-mouse IgG H&L (HRP) at 1:3000 dilution was used as the secondary antibody. Lysate at 20 μ g.




Western blot analysis of lysate from SOX2 protein, using SOX2 Antibody (Cat. #AM2048a). AM2048a was diluted at 1:4000. A goat anti-mouse IgG H&L (HRP) at 1:3000 dilution was used as the secondary antibody. Lysate at 20 μ g.



Formalin-fixed and paraffin-embedded human lung carcinoma tissue reacted with SOX2 Antibody (Cat. #AM2048a), which was

peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Flow cytometric analysis of NCI-H460 cells using SOX2 Monoclonal Antibody (bottom histogram) compared to a negative control cell (top histogram). PE-conjugated goat-anti-mouse secondary antibodies were used for the analysis.

SOX2 Antibody - Background

This intronless gene encodes a member of the SRY-related HMG-box (SOX) family of transcription factors involved in the regulation of embryonic development and in the determination of cell fate. The product of this gene is required for stem-cell maintenance in the central nervous system, and also regulates gene expression in the stomach. Mutations in this gene have been associated with optic nerve hypoplasia and with syndromic microphthalmia, a severe form of structural eye malformation. This gene lies within an intron of another gene called SOX2 overlapping transcript (SOX2OT).

SOX2 Antibody - References

References for protein:

1. Effects of MCF2L2, ADIPOQ and SOX2 genetic polymorphisms on the development of nephropathy in type 1 Diabetes Mellitus. Zhang D, et al. BMC Med Genet, 2010 Jul 28. PMID 20667095.
2. Sox2 protein expression is an independent poor prognostic indicator in stage I lung adenocarcinoma. Sholl LM, et al. Am J Surg Pathol, 2010 Aug. PMID 20631605.
3. Sox2 is translationally activated by eukaryotic initiation factor 4E in human glioma-initiating cells. Ge Y, et al. Biochem Biophys Res Commun, 2010 Jul 9. PMID 20537983.
4. Mutational screening of CHX10, GDF6, OTX2, RAX and SOX2 genes in 50 unrelated microphthalmia-anophthalmia-coloboma (MAC) spectrum cases. Gonzalez-Rodriguez J, et al. Br J Ophthalmol, 2010 Aug. PMID 20494911.
4. Examination of SOX2 in variable ocular conditions identifies a recurrent deletion in

microphthalmia and lack of mutations in other phenotypes. Reis LM, et al. Mol Vis, 2010 Apr 28. PMID 20454695.

References for SY5Y (SH-SY5Y; ATCC#CRL-2266):
1. Ross RA, et al. Coordinate morphological and biochemical interconversion of human neuroblastoma cells. J. Natl. Cancer Inst. 71: 741-749, 1983. [PubMed: 6137586]; 2. Biedler JL, et al. Multiple neurotransmitter synthesis by human neuroblastoma cell lines and clones. Cancer Res. 38: 3751-3757, 1978. [PubMed: 29704].

References for MCF7 cell line:

1. Soule HD; Vazquez J; Long A; Albert S; Brennan M. (1973). "A human cell line from a pleural effusion derived from a breast carcinoma". Journal of the National Cancer Institute 51 (5): 1409-1416. [PMID 4357757].

2. Levenson, AS; Jordan VC. (1997). "MCF-7: the first hormone-responsive breast cancer cell line". Cancer Research 57 (15): 3071-3078. [PMID 9242427].

3. Lacroix, M; Leclercq G. (2004). "Relevance of breast cancer cell lines as models for breast tumours: an update". Breast Research and Treatment 83 (3): 249-289.[PMID 14758095].

SOX2 Antibody - Citations

- [CRISPR/Cas9 targeted deletion of polyglutamine in spinocerebellar ataxia type 3 derived induced pluripotent stem cells.](#)
- [Physiological genomics identifies genetic modifiers of long QT syndrome type 2 severity.](#)
- [Generation of GZKHOi001-A and GZWWTi001-A, two induced pluripotent stem cell lines derived from peripheral blood mononuclear cells of Duchenne muscular dystrophy patients.](#)
- [Generation of integration-free induced pluripotent stem cells \(GZHMUi001-A\) by reprogramming peripheral blood mononuclear cells from a 47, XXX syndrome patient.](#)
- [Brother of the regulator of the imprinted site \(BORIS\) variant subfamily 6 is a novel target of lung cancer stem-like cell immunotherapy.](#)
- [Melatonin Inhibits Glioblastoma Stem-like cells through Suppression of EZH2-NOTCH1 Signaling Axis.](#)
- [ROCK Inhibition Facilitates In Vitro Expansion of Glioblastoma Stem-Like Cells.](#)